arXiv:1105.5770v2 [math.CA] 20 Sep 2012

A connection formula of the q-confluent hypergeometric function

Takeshi MORITA[*]

Abstract

We show a connection formula of the q-confluent hypergeometric function. We introduce the q-Borel and q-Laplace transformations following C. Zhang to obtain the connection formula. Combining with Zhang's formula, we show the connection matrix. We give the limit $q \to 1-0$ of our connection formula.

1 Introduction

We show a new connection formula of solutions to the q-confluent hypergeometric equation

$$(1-abqx)u(q^2x)-\{1-(a+b)qx\}\,u(qx)-qxu(x)=0. \tag{1}$$

The equation (1) has solutions

$$u_1(x)={}_2\varphi_0(a,b;-;q,x), \quad u_2(x)=\frac{(abx;q)_\infty}{\theta(-qx)}{}_2\varphi_1\left(\frac{q}{a},\frac{q}{b};0;q,abx\right)$$

around the origin and has solutions

$$v_1(x)=x^{-\alpha}{}_2\varphi_1\left(a,0;\frac{aq}{b};q,\frac{q}{abx}\right), \quad v_2(x)=x^{-\beta}{}_2\varphi_1\left(b,0;\frac{bq}{a};q,\frac{q}{abx}\right)$$

---

[*] Graduate School of Information Science and Technology, Osaka University, 1-1 Machikaneyama-machi, Toyonaka, 560-0043, Japan. E-mail: t-morita@cr.math.sci.osaka-u.ac.jp

around the infinity. Here $q^\alpha = a$ and $q^\beta = b$.

The connection formula of a q-difference linear equation of the second order is a linear relation between $u_1(x), u_2(x)$ and $v_1(x), v_2(x)$:

$$\begin{pmatrix} u_1(x) \\ u_2(x) \end{pmatrix} = \begin{pmatrix} C_{11}(x) & C_{12}(x) \\ C_{21}(x) & C_{22}(x) \end{pmatrix} \begin{pmatrix} v_1(x) \\ v_2(x) \end{pmatrix},$$

where the connection coefficient $C_{jk}(x)$ is a q-periodic function.

Zhang has shown a connection formula for $u_1(x)$ [9]. We show a connection formula for $u_2(x)$:

$$\begin{aligned} {}_2\varphi_1(q/a, q/b; 0; q, abx) = & \frac{(q/a;q)_\infty}{(b/a;q)_\infty} \frac{(aqx, 1/ax; q)_\infty}{(abx;q)_\infty} {}_2\varphi_1(a, 0; aq/b; q, 1/abx) \\ & + \frac{(q/b;q)_\infty}{(a/b;q)_\infty} \frac{(bqx, 1/bx; q)_\infty}{(abx;q)_\infty} {}_2\varphi_1(b, 0; bq/a; q, 1/abx). \end{aligned}$$

Since $u_1(x)$ is a divergent series, the q-Stokes phenomenon appears in Zhang's connection formula. But our connection formula does not have any discontinuity.

We fix our notations, following by [2]. The function $(a;q)_n$ is the q-shifted factorial:

$$(a;q)_n := \begin{cases} 1, & n = 0, \\ (1-a)(1-aq)\cdots(1-aq^{n-1}), & n \geq 1, \end{cases}$$

$$(a;q)_\infty = \lim_{n\to\infty} (a;q)_n$$

and

$$(a_1, a_2, \cdots, a_m; q)_\infty = (a_1;q)_\infty (a_2;q)_\infty \cdots (a_m;q)_\infty.$$

Moreover, the basic hypergeometric series ${}_r\varphi_s$ is

$${}_r\varphi_s(a_1, \cdots, a_r; b_1, \cdots, b_s; q, x) := \sum_{n\geq 0} \frac{(a_1, \cdots, a_r; q)_n}{(b_1, \cdots, b_s; q)_n (q;q)_n} \left[(-1)^n q^{\frac{n(n-1)}{2}}\right]^{1+s-r} x^n.$$

We assume that $q \in \mathbb{C}^*$ satisfies $0 < |q| < 1$. The q-difference operator σ_q is given by $\sigma_q f(x) = f(qx)$. The theta function of Jacobi is given by following series:

$$\theta_q(x) := \sum_{n\in\mathbb{Z}} q^{\frac{n(n-1)}{2}} x^n, \quad \forall x \in \mathbb{C}^*.$$

We denote by $\theta_q(x)$ or more shortly $\theta(x)$. Jacobi's triple product identity is

$$\theta(x) = \left(q, -x, -\frac{q}{x}; q\right)_\infty .$$

The theta function has inversion formula $\theta(x) = x\theta(1/x)$. For all fixed $\lambda \in \mathbb{C}^*$, we define a q-spiral $[\lambda; q] := \lambda q^{\mathbb{Z}} = \{\lambda q^k; k \in \mathbb{Z}\}$. We remark that $\theta(\lambda q^k/x) = 0$ if and only if $x \in [-\lambda; q]$.

We study connection problems on linear q-difference equations with irregular singular points. Connection problems on linear q-difference equations of the Laplace type [5] with regular singular points are studied by G. D. Birkhoff [1]. Generically, linear q-difference equations has formal power series solutions $x^\alpha \sum_{n\geq 0} a_n x^n$ around the origin and $x^\beta \sum_{n\geq 0} b_n x^{-n}$ around the infinity. But on connection problems on linear q-difference equations, we replace the function x^κ with the function $\theta(x)/\theta(kx)$, where $k = q^\kappa$. Then, the fundamental system of solutions are given by single valued functions which have single poles at suitable q-spirals. The first example is given by G. N. Watson [6]. But a few examples of irregular singular case are known [7, 8, 4]. In this paper, we give a connection formula of the q-confluent type function with using q-Borel-Laplace transformations.

At first, we review the confluent hypergeometric equation (CHGE). In 1812, C. F. Gauss introduced the hypergeometric series ${}_2F_1(\alpha, \beta; \gamma; z)$ as follows:

$$ {}_2F_1(\alpha, \beta; \gamma; z) = \sum_{n\geq 0} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n n!} z^n, \quad \gamma \neq 0, -1, -2, \cdots . \tag{2}$$

Here,

$$(\alpha)_n = \alpha\{\alpha + 1\} \cdots \{\alpha + (n-1)\}.$$

More generally, the generalized hypergeometric series is given by

$$ {}_rF_s(\alpha_1, \cdots, \alpha_r; \beta_1, \cdots, \beta_s; z) = \sum_{n\geq 0} \frac{(\alpha_1)_n \cdots (\alpha_r)_n}{(\beta_1)_n \cdots (\beta_s)_n n!} z^n.$$

The hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$ satisfies the second order differential equation

$$z(1-z)\frac{d^2u}{dz^2} + \{\gamma - (\alpha + \beta + 1)z\}\frac{du}{dz} - \alpha\beta u = 0. \tag{3}$$

Gauss gives the connection formula of the function ${}_2F_1(\alpha,\beta;\gamma,z)$. We put $z \mapsto z/\beta$ and take the limit $\beta \to \infty$ in the equation (3), we obtain the confluent hypergeometric equation (CHGE)

$$z\frac{d^2u}{dz^2} + (\gamma - z)\frac{du}{dz} - \alpha u = 0. \tag{4}$$

Solutions of (4) around the origin are

$$u_1(z) = {}_1F_1(\alpha;\gamma;z) \tag{5}$$

and

$$u_2(z) = z^{1-\gamma}{}_1F_1(\alpha-\gamma+1, 2-\gamma, z). \tag{6}$$

Solutions around the infinity are given by divergent series

$$v_1(z) = (-z)^{-\alpha}{}_2F_0(\alpha, \alpha-\gamma+1; -; 1/z) \tag{7}$$

and

$$v_2(z) = e^z z^{\alpha-\gamma}{}_2F_0(1-\alpha, \gamma-\alpha; -; 1/z). \tag{8}$$

A well-known connection formula between (7), (8) and (5) is given by

$$\begin{aligned}{}_1F_1(\alpha;\gamma;z) \sim& \frac{\Gamma(\gamma)}{\Gamma(\gamma-\alpha)}(-z)^{-\alpha}{}_2F_0(\alpha,\alpha-\gamma+1;-;-1/z)\\ &+\frac{\Gamma(\gamma)}{\Gamma(\alpha)}e^z z^{\alpha-\gamma}{}_2F_0(1-\alpha,\gamma-\alpha;-;1/z)\end{aligned}$$

where $-\pi/2 < \arg z < 3\pi/2$. We remark that the connection formula for second solution around the infinity (6) can be derived from the first one. In section two, we deal with the degenaration of the equation (3) which is slightly different from standard way.

It is known that there exists the q-analogue of ${}_2F_1(\alpha,\beta;\gamma;z)$, which is introduced by E. Heine in 1847 as follows:

$${}_2\varphi_1(a,b;c;q,x) := \sum_{n\geq 0}\frac{(a;q)_n(b;q)_n}{(c;q)_n(q;q)_n}x^n.$$

The function ${}_2\varphi_1(a,b;c;q,x)$ satisfies the second order q-difference equation (q-HGE)

$$\begin{aligned}x(c-abqx)\mathcal{D}_q^2u + \left[\frac{1-c}{1-q} + \frac{(1-a)(1-b)-(1-abq)}{1-q}x\right]\mathcal{D}_q u \\ - \frac{(1-a)(1-b)}{(1-q)^2}u = 0, \quad |x| < 1,\end{aligned} \tag{9}$$

where $\mathcal{D}_q$ is the q-derivative operator defined for fixed q by

$$\mathcal{D}_q f(x) = \frac{f(x) - f(qx)}{(1-q)x}.$$

By replacing a, b, c by q^α, q^β, q^γ and then letting $q \to 1-0$, the equation (9) tends to the hypergeometric equation (3) where $|x| < 1$. The q-hypergeometric equation (9) can be rewritten as follows:

$$(c - abqx)u(q^2x) - \{c + q - (a+b)qx\}\, u(qx) + q(1-x)u(x) = 0. \tag{10}$$

If we set $x \mapsto cx$ and $c \to \infty$ in (10), we obtain the q-confluent hypergeometric equation (1). The equation (1) is considered as a q-analogue of CHGE. We remark that the first solution $u_1(x)$ is a divergent series and $u_2(x)$ is a convergent around the origin. The second solution $u_2(x)$ can not be derived from $u_1(x)$ directly. This point is essentially different from the differential equation case.

Connection problems on linear q-difference equations between the origin and the infinity are studied by G. D. Birkhoff [1]. In 1910, Watson [6] has shown the connection formula of the series ${}_2\varphi_1(a, b; c; q, x)$ as follows:

$$\begin{aligned} {}_2\varphi_1(a, b; c; q; x) = {} & \frac{(b, c/a; q)_\infty (ax, q/ax; q)_\infty}{(c, b/a; q)_\infty (x, q/x; q)_\infty} {}_2\varphi_1(a, aq/c; aq/b; q, cq/abx) \\ & + \frac{(a, c/b; q)_\infty (bx, q/bx; q)_\infty}{(c, a/b; q)_\infty (x, q/x; q)_\infty} {}_2\varphi_1(b, bq/c; bq/a; q, cq/abx). \end{aligned}$$

We remark that we can not set $c = 0$ directly in this formula.

In 2002, C. Zhang [9] shows one of the connection formula of the q-CHGE as follows:

$$\begin{aligned} {}_2f_0(a, b; \lambda, q, x) = {} & \frac{(b; q)_\infty}{\left(\frac{b}{a}; q\right)_\infty} \frac{\theta(a\lambda)}{\theta(\lambda)} \frac{\theta\left(\frac{qax}{\lambda}\right)}{\theta\left(\frac{qx}{\lambda}\right)} {}_2\varphi_1\left(a, 0; \frac{aq}{b}; q, \frac{q}{abx}\right) \\ & + \frac{(a; q)_\infty}{\left(\frac{a}{b}; q\right)_\infty} \frac{\theta(b\lambda)}{\theta(\lambda)} \frac{\theta\left(\frac{qbx}{\lambda}\right)}{\theta\left(\frac{qx}{\lambda}\right)} {}_2\varphi_1\left(b, 0; \frac{bq}{a}; q, \frac{q}{abx}\right). \end{aligned} \tag{11}$$

The function ${}_2f_0(a, b; \lambda, q, x)$ is the q-Borel-Laplace transform of ${}_2\varphi_0(a, b; -; q, x)$, i.e.,

$${}_2f_0(a, b; \lambda, q, x) := \mathcal{L}^+_{q,\lambda} \circ \mathcal{B}^+_q\, {}_2\varphi_0(a, b; -; q, x).$$

The q-Borel-Laplace transformations are studied by C. Zhang in [9] (see section two for more details). When we study connection problems of divergent series, this resummation method becomes a powerful tool. We remark that we can find a new parameter λ in the resummation ${}_2f_0(a, b; \lambda, q, x)$. This parameter brings us new viewpoints for the study of the q-Stokes phenomenon.

It is known that there exists two different types of the q-Borel transformation and the q-Laplace transformation. The q-Borel-Laplace transformation of the first kind are defined in [9] and the q-Borel-Laplace transformations of the second kind are studied in [8]. These q-Borel transformations are the formal inverse transformation of each of the q-Laplace transformations.

Zhang has shown a connection formula of the series ${}_2\varphi_0(a, b; -; q, x)$ by the q-Borel-Laplace transformations of the first kind. But the connection formula for the second solution of (1) has not known. In this paper, we show the second connection formula for q-CHGE with using the q-Borel transformation and the q-Laplace transformation of the second kind. Combining with Zhang's connection formula, we obtain the following connection in section two:

Theorem. *For any* $\lambda, \mu \in \mathbb{C}^*$, $x \in \mathbb{C}^* \setminus [1; q] \cup [-\mu/a; q] \cup [-\lambda; q]$, *we have*

$$\begin{pmatrix} {}_2f_0(a, b; \lambda, q, x) \\ {}_2f_1(a, b; q, x) \end{pmatrix} = \begin{pmatrix} C_\mu^\lambda(a, b; q, x) & C_\mu^\lambda(b, a; q, x) \\ C_\mu(a, b; q, x) & C_\mu(b, a; q, x) \end{pmatrix} \begin{pmatrix} S_\mu(a, b; q, x) \\ S_\mu(b, a; q, x) \end{pmatrix}.$$

The set $[\lambda; q]$ is the q-spiral. The function ${}_2f_1(a, b; q, x)$ is the q-Borel-Laplace transform of ${}_2\varphi_1(a, b; 0; q, x)$. The function $S_\mu(a, b; q, x)$ is the solution of (1) around the infinity as follows;

$$S_\mu(a, b; q, x) := \frac{\theta(a\mu x)}{\theta(\mu x)} {}_2\varphi_1\left(a, 0; \frac{aq}{b}; q, \frac{q}{abx}\right).$$

We remark that the function $x^{-\alpha}$ in the solution $v_1(x)$ is replaced with the function $\theta(a\mu x)/\theta(\mu x)$, where $a = q^\alpha$. Functions $C_\mu^\lambda(a, b; q, x)$, $C_\mu(a, b; q, x)$ are q-elliptic functions.

In section three, we consider the limit $q \to 1-0$ of our connection formula. If we take a limit $q \to 1 - 0$ of our connection formula, we formally obtain the connection formula of the confluent hypergeometric series ${}_2F_0$.

2 The connection formula and the connection matrix

We review a q-confluent hypergeometric equation in subsection 2.1. Then we show a connection formula for the q-confluent hypergeometric function, which is different from Zhang's formula.

2.1 Confluent hypergeometric equation

For the confluent hypergeometric equation (3), we take another degeneration. We put $z \mapsto z\gamma$ and the take the limit $\gamma \to 0$. Then we obtain

$$z^2 \frac{d^2u}{dz^2} - \{1-(\alpha+\beta+1)z\}\frac{du}{dz} + \alpha\beta u = 0. \tag{12}$$

Solutions of (12) around the origin are given by divergent series

$$\tilde{u}_1(z) = {}_2F_0(\alpha, \beta; -, z)$$

and

$$\tilde{u}_2(z) = e^{\frac{1}{z}}(-z)^{1-\alpha-\beta}{}_2F_0(1-\alpha, 1-\beta; -, z).$$

Solutions around the infinity are

$$\tilde{v}_1(z) = (-z)^{\alpha}{}_1F_1(\alpha, 1+\alpha-\beta, -1/z)$$

and

$$\tilde{v}_2(z) = (-z)^{\beta}{}_1F_1(\beta, 1+\beta-\alpha, -1/z).$$

We consider the following second order q-difference equation

$$x(-1+abqx)D_q^2u(x) + \left\{\frac{1}{1-q} - \frac{(1-a)(1-b)-(1-abq)}{1-q}x\right\}D_qu(x) + \frac{(1-a)(1-b)}{(1-q)^2}u(x) = 0.$$

This equation can be rewritten as follows:

$$(1-abqx)u(xq^2) - \{1-(a+b)qx\}u(xq) - qxu(x) = 0, \tag{13}$$

which is called a q-confluent hypergeometric equation.

2.2 Local solutions to the q-confluent hypergeometric equation

In the following, we study connection problem of (13). At first we show local solutions to (13) around $x = 0$ and $x = \infty$.

Lemma 1. *The equation (13) has solutions*

$$u_1(x) = {}_2\varphi_0(a, b; -; q, x), \tag{14}$$

$$u_2(x) = \frac{(abx; q)_\infty}{\theta(-qx)} {}_2\varphi_1 \left(\frac{q}{a}, \frac{q}{b}; 0; q, abx \right) \tag{15}$$

around the origin and has solutions

$$v_1(x) = x^{-\alpha} {}_2\varphi_1 \left(a, 0; \frac{aq}{b}; q, \frac{q}{abx} \right), \tag{16}$$

$$v_2(x) = x^{-\beta} {}_2\varphi_1 \left(b, 0; \frac{bq}{a}; q, \frac{q}{abx} \right), \tag{17}$$

around the infinity, provided that $a = q^\alpha$ and $b = q^\beta$.

Proof. We show a fundamental system of solutions of (13) around $x = 0$. We set $u(x) = \sum_{n \geq 0} a_n x^n$, $a_0 = 1$. Then we obtain

$$u_1(x) = {}_2\varphi_0(a, b; -; q, x).$$

We find another solution. We set $\mathcal{E}(x) = 1/\theta(-qx)$ and $f(x) = \sum_{n \geq 0} a_n x^n$, $a_0 = 1$ to obtain the solution around the origin. We assume that $u(x) = \mathcal{E}(x) f(x)$. We remark that the function $\mathcal{E}(x)$ has following property:

$$\sigma_q \mathcal{E}(x) = -qx\mathcal{E}(x), \qquad \sigma_q^2 \mathcal{E}(x) = q^3 x^2 \mathcal{E}(x).$$

Therefore, we obtain the equation

$$\left[q^3 x (1 - abqx) \sigma_q^2 + q \left\{ 1 - (a + b) qx \right\} \sigma_q - q \right] f(x) = 0. \tag{18}$$

Since the infinite product $(abx; q)_\infty$ satisfies the following q-difference relation

$$\sigma_q (abx; q)_\infty = \frac{1}{1 - abx} (abx; q)_\infty,$$

we obtain the second solution. Therefore, solutions of the equation (13) are given by

$$u_1(x) = {}_2\varphi_0(a,b;-;q,x), \quad u_2(x) = \frac{(abx;q)_\infty}{\theta(-qx)} {}_2\varphi_1\left(\frac{q}{a},\frac{q}{b};0;q,abx\right).$$

Around $x=\infty$, we can easily determine local solutions by setting

$$v(x) = \frac{\theta(a\mu x)}{\theta(\mu x)} \sum_{n\geq 0} a_n x^{-n}, \quad a_0 = 1,$$

for any fixed $\mu \in \mathbb{C}^*$ and $x \in \mathbb{C}^* \setminus [-\mu;q]$. □

Here, $u_1(x)$ is a divergent series and $u_2(x)$,$v_1(x)$ and $v_2(x)$ are convergent series [2]. Therefore, the q-Stokes phenomenon occurs for $u_1(x)$ and Zhang has shown a connection formula for $u_1(x)$ be means of the following q-Borel-Laplace transformations of the first kind [9].

Definition 1. *For any* $f(x) = \sum_{n\geq 0} a_n x^n$*, The q-Borel transformation* $\mathcal{B}_q^+$ *is*

$$\left(\mathcal{B}_q^+ f\right)(\xi) = \varphi(\xi) := \sum_{n\geq 0} a_n q^{\frac{n(n-1)}{2}} \xi^n,$$

the q-Laplace transformation $\mathcal{L}_{q,\lambda}^+$ *is*

$$\left(\mathcal{L}_{q,\lambda}^+ \varphi\right)(x) := \sum_{n\in\mathbb{Z}} \frac{\varphi(q^n\lambda)}{\theta\left(\frac{q^n\lambda}{x}\right)}.$$

C. Zhang has determined a resummation of (14) by the q-Borel-Laplace transformations of the first kind as follows:

$${}_2f_0(a,b;\lambda,q,x) := \mathcal{L}_{q,\lambda}^+ \circ \mathcal{B}_q^+ {}_2\varphi_0(a,b;-;q,x).$$

Then he has given a connection formula for ${}_2f_0(a,b;\lambda,q,x)$.

$$\begin{aligned} {}_2f_0(a,b;\lambda,q,x) = & \frac{(b;q)_\infty}{\left(\frac{b}{a};q\right)_\infty} \frac{\theta(a\lambda)}{\theta(\lambda)} \frac{\theta\left(\frac{qax}{\lambda}\right)}{\theta\left(\frac{qx}{\lambda}\right)} {}_2\varphi_1\left(a,0;\frac{aq}{b};q,\frac{q}{abx}\right) \\ & + \frac{(a;q)_\infty}{\left(\frac{a}{b};q\right)_\infty} \frac{\theta(b\lambda)}{\theta(\lambda)} \frac{\theta\left(\frac{qbx}{\lambda}\right)}{\theta\left(\frac{qx}{\lambda}\right)} {}_2\varphi_1\left(b,0;\frac{bq}{a};q,\frac{q}{abx}\right), \end{aligned}$$

for $x \in \mathbb{C}^* \setminus [-\lambda;q]$.

But the connection formula between (15) and (16),(17) is not known. In the next section, we show the second connection formula with using the q-Borel-Laplace transformations of the second kind.

2.3 The second connection formula

We introduce the q-Borel transformation and the q-Laplace transformation of the second kind following C. Zhang to obtain the solution of the equation (18).

Definition 2. *For $f(x) = \sum_{n\geq 0} a_n t^n$, the q-Borel transformation is defined by*

$$g(\xi) = \left(\mathcal{B}_q^- f\right)(\xi) := \sum_{n\geq 0} a_n q^{-\frac{n(n-1)}{2}} \xi^n,$$

and the q-Laplace transformation is given by

$$\left(\mathcal{L}_q^- g\right)(x) := \frac{1}{2\pi i} \int_{|\xi|=r} g(\xi)\theta_q\left(\frac{x}{\xi}\right)\frac{d\xi}{\xi}.$$

Here, $r_0 > 0$ is enough small number. The q-Borel transformation is considered as a formal inverse of the q-Laplace transformation.

Lemma 2. *We assume that the function f can be q-Borel transformed to the analytic function $g(\xi)$ around $\xi = 0$. Then, we have*

$$\mathcal{L}_q^- \circ \mathcal{B}_q^- f = f.$$

Proof. We can show this lemma calculating residues of the q-Laplace transformation around the origin. ☐

The q-Borel transformation satisfies the following operational relation.

Lemma 3. *For any $l, m \in \mathbb{Z}_{\geq 0}$,*

$$\mathcal{B}_q^-\left(x^m \sigma_q^l\right) = q^{-\frac{m(m-1)}{2}} \xi^m \sigma_q^{l-m} \mathcal{B}_q^-.$$

We apply the q-Borel transformation to the equation (18) and using Lemma 3. We check out $g(\xi)$ satisfies the first order difference equation

$$g(q\xi) = \frac{(1+aq\xi)(1+bq\xi)}{(1+q^2\xi)} g(\xi).$$

Since $g(0) = a_0 = 1$, we have the infinite product of $g(\xi)$ as follows;

$$g(\xi) = \frac{(-q^2\xi;q)_\infty}{(-qa\xi;q)_\infty(-qb\xi;q)_\infty}.$$

We remark that $g(\xi)$ has single poles at

$$\left\{\xi \in \mathbb{C}^*; \xi = -\frac{1}{aq^{k+1}}, -\frac{1}{bq^{k+1}}, k \in \mathbb{Z}_{\geq 0}\right\}.$$

We set

$$0 < r < r_0 := \max\left\{\frac{1}{|aq|}, \quad \frac{1}{|bq|}\right\}$$

and choose the radius $r > 0$ such that $0 < r < r_0$. By Cauchy's residue theorem, the p-Laplace transform of $g(\xi)$ is

$$\begin{aligned} f(x) =& \frac{1}{2\pi i}\int_{|\xi|=r} g(\xi)\theta\left(\frac{x}{\xi}\right)\frac{d\xi}{\xi} \\ =& -\sum_{k\geq 0} \mathrm{Res}\left\{g(\xi)\theta\left(\frac{x}{\xi}\right)\frac{1}{\xi}; \xi = -\frac{1}{aq^{k+1}}\right\} \\ & -\sum_{k\geq 0} \mathrm{Res}\left\{g(\xi)\theta\left(\frac{x}{\xi}\right)\frac{1}{\xi}; \xi = -\frac{1}{bq^{k+1}}\right\} \end{aligned}$$

where $0 < r < r_0$. We use the following lemma to calculate the residue.

Lemma 4. *For any $k \in \mathbb{N}$, $\lambda \in \mathbb{C}^*$, we have;*

1. $\mathrm{Res}\left\{\frac{1}{(\xi/\lambda; q)_\infty}\frac{1}{\xi} : \xi = \lambda q^{-k}\right\} = \frac{(-1)^{k+1} q^{\frac{k(k+1)}{2}}}{(q;q)_k (q;q)_\infty}$,

2. $\frac{1}{(\lambda q^{-k}; q)_\infty} = \frac{(-\lambda)^{-k} q^{\frac{k(k+1)}{2}}}{(\lambda; q)_\infty \left(q/\lambda; q\right)_k}, \quad \lambda \notin q^{\mathbb{Z}}$.

Summing up all of residues, we obtain $f(x)$ as follows:

$$\begin{aligned} f(x) =& \frac{\left(\frac{q}{a}; q\right)_\infty}{\left(\frac{b}{a}, q; q\right)_\infty}\frac{\theta(-aqx)}{\theta(-qx)}\,{}_2\varphi_1\left(a, 0; \frac{aq}{b}; q, \frac{q}{abx}\right) \\ & + \frac{\left(\frac{q}{b}; q\right)_\infty}{\left(\frac{a}{b}, q; q\right)_\infty}\frac{\theta(-bqx)}{\theta(-qx)}\,{}_2\varphi_1\left(b, 0; \frac{bq}{a}; q, \frac{q}{abx}\right). \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 1. *For any $x \notin [1;q]$, we have*

$$\begin{aligned}u_2(x) =&\frac{(abx;q)_\infty}{\theta(-qx)}{}_2\varphi_1\left(\frac{q}{a},\frac{q}{b};0;q,abx\right)\\ =&\frac{\left(\frac{q}{a};q\right)_\infty}{\left(\frac{b}{a},q;q\right)_\infty}\frac{\theta(-aqx)}{\theta(-qx)}{}_2\varphi_1\left(a,0;\frac{aq}{b};q,\frac{q}{abx}\right)\\ &+\frac{\left(\frac{q}{b};q\right)_\infty}{\left(\frac{a}{b},q;q\right)_\infty}\frac{\theta(-bqx)}{\theta(-qx)}{}_2\varphi_1\left(b,0;\frac{bq}{a};q,\frac{q}{abx}\right).\end{aligned}$$

2.4 The connection matrix

We give the connection matrix for the equation (1). We define new functions $S_\mu(a,b;q,x)$, $C_\mu^\lambda(a,b;q,x)$ and $C_\mu(a,b;q,x)$. For any λ, $\mu \in \mathbb{C}^*$, $x \in \mathbb{C}^* \setminus [-\mu;q]$, $S_\mu(a,b;q,x)$ is

$$S_\mu(a,b;q,x) := \frac{\theta(a\mu x)}{\theta(\mu x)}{}_2\varphi_1\left(a,0;\frac{aq}{b};q,\frac{q}{abx}\right).$$

The function $\theta(a\mu x)/\theta(\mu x)$ satisfies the following q-difference equation

$$u(qx) = \frac{1}{a}u(x)$$

which is also satisfied by the function $u(x) = x^{-\alpha}$, $a = q^\alpha$. We remark that the pair $(S_\mu(a,b;q,x), S_\mu(b,a;q,x))$ gives a fundamental system of solutions of the equation (1) if $a, b \notin \mathbb{Z}$. We set functions $C_\mu^\lambda(a,b;q,x)$ and $C_\mu(a,b;q,x)$ as follow;

Definition 3. *For any $\lambda, \mu \in \mathbb{C}^*$, $C_\mu^\lambda(a,b;q,x)$ is*

$$C_\mu^\lambda(a,b;q,x) := \frac{(b;q)_\infty}{\left(\frac{b}{a};q\right)_\infty}\frac{\theta(a\lambda)}{\theta(\lambda)}\frac{\theta\left(\frac{qax}{\lambda}\right)}{\theta\left(\frac{qx}{\lambda}\right)}\frac{\theta(\mu x)}{\theta(a\mu x)}.$$

Similarly, $C_\mu(a,b;q,x)$ is

$$C_\mu(a,b;q,x) := \frac{\left(\frac{q}{a};q\right)_\infty}{\left(\frac{b}{a},q;q\right)}\frac{\theta(-aqx)}{\theta(-qx)}\frac{\theta(\mu x)}{\theta(a\mu x)}.$$

Then $C_\mu^\lambda(a,b;q,x)$ and $C_\mu(a,b;q,x)$ are single valued as a function of x. They satisfy the following relation

$$C_\mu^\lambda(a,b;q,e^{2\pi i}x)=C_\mu^\lambda(a,b;q,x),\quad C_\mu^\lambda(a,b;q,qx)=C_\mu^\lambda(a,b;q,x)$$

and

$$C_\mu(a,b;q,e^{2\pi i}x)=C_\mu(a,b;q,x),\quad C_\mu(a,b;q,qx)=C_\mu(a,b;q,x).$$

i.e. $C_\mu^\lambda(a,b;q,x)$ and $C_\mu(a,b;q,x)$ are q-elliptic functions. We set

$${}_2f_1(a,b;q,x):=u_2(x)=\frac{(abx;q)_\infty}{\theta(-qx)}{}_2\varphi_1\left(\frac{q}{a},\frac{q}{b};0;q,abx\right).$$

Combining Zhang's formula (11) and Theorem 1, we obtain the connection formula in matrix form.

Theorem 2. *For any $\lambda,\mu\in\mathbb{C}^*$, $x\in\mathbb{C}^*\setminus[1;q]\cup[-\mu/a;q]\cup[-\lambda;q]$, we have*

$$\begin{pmatrix}{}_2f_0(a,b;\lambda,q,x)\\ {}_2f_1(a,b;q,x)\end{pmatrix}=\begin{pmatrix}C_\mu^\lambda(a,b;q,x) & C_\mu^\lambda(b,a;q,x)\\ C_\mu(a,b;q,x) & C_\mu(b,a;q,x)\end{pmatrix}\begin{pmatrix}S_\mu(a,b;q,x)\\ S_\mu(b,a;q,x)\end{pmatrix}.$$

3 The limit $q\to 1-0$ of the connection formula

In this section, we show the limit $q\to 1-0$ of our connection formula. In [9], Zhang has given the limit $q\to 1-0$ of as follows:

Theorem 3 (Zhang)**.** *For any $\alpha,\beta\in\mathbb{C}^*(\alpha-\beta\notin\mathbb{Z})$ and z in any compact domain of $\mathbb{C}^*\setminus[-\infty,0]$, we have*

$$\begin{aligned}\lim_{q\to 1-0}{}_2f_0\left(q^\alpha,q^\beta;\lambda,q,\frac{z}{(1-q)}\right)&=\frac{\Gamma(\beta-\alpha)}{\Gamma(\beta)}z^{-\alpha}{}_1F_1\left(\alpha,\alpha-\beta+1;\frac{1}{z}\right)\\ &\quad+\frac{\Gamma(\alpha-\beta)}{\Gamma(\alpha)}z^{-\beta}{}_1F_1\left(\beta;\beta-\alpha+1;\frac{1}{z}\right).\end{aligned}$$

Our limit of the connection formula in theorem 1 is different from the theorem above. The aim of this section is to give the following theorem.

Theorem 4. *The limit $q \to 1-0$ of our connection formula formally gives the following asymptotic formula;*

$$\begin{aligned} e^{1/z}(-z)^{1-\alpha-\beta}{}_2F_0(1-\alpha,1-\beta;-,z) &= \frac{\Gamma(\beta-\alpha)}{\Gamma(1-\alpha)}(-z)^{-\alpha}{}_1F_1\left(\alpha;\alpha+1-\beta;\frac{1}{z}\right) \\ &+\frac{\Gamma(\alpha-\beta)}{\Gamma(1-\beta)}(-z)^{-\beta}{}_1F_1\left(\beta;\beta+1-\alpha;\frac{1}{z}\right). \end{aligned}$$

We review some limits. According to [7], we have the following proposition. We take principal value of the logarithm on $\mathbb{C}^* \setminus (-\infty, 0]$.

Proposition 1. *For any $\gamma \in \mathbb{C}^*$, we have*

$$\lim_{q\to 1-0} \frac{\theta\left(q^\gamma \frac{u}{1-q}\right)}{\theta\left(\frac{u}{1-q}\right)}(1-q)^{-\gamma} = u^{-\gamma}$$

converges uniformly on compact subset of $\mathbb{C} \setminus (-\infty, 0]$.

We also review the q-gamma function $\Gamma_q(\cdot)$ and the q-exponential function $E_q(\cdot)$. The q-gamma function is defined by

$$\Gamma_q(x) := \frac{(q;q)_\infty}{(q^x;q)_\infty}(1-q)^{1-x}, \quad 0<q<1.$$

This function satisfies $\lim_{q\to 1-0}\Gamma_q(x) = \Gamma(x)$ [2]. The q-exponential function $E_q(\cdot)$ is

$$E_q(z) = \sum_{n\geq 0} \frac{q^{n(n-1)/2}}{(q;q)_n} z^n = (-z;q)_\infty.$$

The function $E_q(\cdot)$ satisfies the following limit. In this sense, the q-exponential function is considered as a q-analogue of the exponential function:

$$\lim_{q\to 1-0} E_q\left(z(1-q)\right) = e^z.$$

We set $a = q^\alpha$, $b = q^\beta$ and $x = z/(1-q)$ in the theorem 1. We introduce new constant $w(\alpha,\beta;q)$ as follows:

$$w(\alpha,\beta;q) := (q;q)_\infty(1-q)^{1-\alpha-\beta}.$$

We consider each side of the limit $q \to 1-0$ of $w(\alpha,\beta;q){}_2f_1\left(q^\alpha, q^\beta; q, z/(1-q)\right)$.

Lemma 5. *For any* $\alpha, \beta \in \mathbb{C}^*, (\alpha - \beta \notin \mathbb{Z})$*, we have*

$$\lim_{q \to 1-0} w(\alpha, \beta; q) \frac{\left(\frac{q^{\alpha+\beta} z}{1-q}; q\right)_\infty}{\theta\left(-\frac{qz}{1-q}\right)} {}_2\varphi_1\left(q^{1-\alpha}, q^{1-\beta}; 0; q, \frac{q^{\alpha+\beta} z}{1-q}\right)$$
$$= (-z)^{1-\alpha-\beta} e^{\frac{1}{z}} {}_2F_0(1-\alpha, 1-\beta; -, z).$$

Proof. We remark that

$$\begin{aligned} w(\alpha, \beta; q) \frac{\left(\frac{q^{\alpha+\beta} z}{1-q}; q\right)_\infty}{\theta\left(-\frac{qz}{1-q}\right)} {}_2\varphi_1\left(q^{1-\alpha}, q^{1-\beta}; 0; q, \frac{q^{\alpha+\beta} z}{1-q}\right) \\ = \left\{ \frac{\theta\left(q^{\alpha+\beta-1} \frac{-qz}{1-q}\right)}{\theta\left(\frac{-qz}{1-q}\right)} (1-q)^{1-\alpha-\beta} \right\} \frac{1}{E_q\left(-\frac{(1-q)}{q^{\alpha+\beta-1} z}\right)} \\ \times {}_2\varphi_1\left(q^{1-\alpha}, q^{1-\beta}; 0; q, \frac{q^{\alpha+\beta} z}{1-q}\right), \end{aligned}$$

we obtain the conclusion. □

We consider the other hand side.

Lemma 6. *For any* $\alpha, \beta \in \mathbb{C}^*, (\alpha - \beta \notin \mathbb{Z})$*, we have*

$$\begin{aligned} \lim_{q \to 1-0} w(\alpha, \beta, q) {}_2f_1\left(q^\alpha, q^\beta; q, \frac{z}{(1-q)}\right) = & \frac{\Gamma(\beta-\alpha)}{\Gamma(1-\alpha)} (-z)^{-\alpha} {}_1F_1\left(\alpha; \alpha+1-\beta; \frac{1}{z}\right) \\ & + \frac{\Gamma(\alpha-\beta)}{\Gamma(1-\beta)} (-z)^{-\beta} {}_1F_1\left(\beta; \beta+1-\alpha; \frac{1}{z}\right). \end{aligned}$$

Proof. We remark that

$$\begin{aligned} w(\alpha, \beta; q) \frac{(q^{1-\alpha}; q)_\infty}{(q^{\beta-\alpha}, q; q)_\infty} \frac{\theta\left(-\frac{q^{\alpha+1} z}{1-q}\right)}{\theta\left(-\frac{qz}{1-q}\right)} {}_2\varphi_1\left(q^\alpha, 0; q^{\alpha+1-\beta}; q, \frac{q^{1-\alpha-\beta}(1-q)}{z}\right) \\ = \left\{ \frac{(q^{1-\alpha}; q)_\infty}{(q; q)_\infty} (1-q)^\alpha \right\} \left\{ \frac{(q; q)_\infty}{(q^{\beta-\alpha}; q)_\infty} (1-q)^{-(1-\beta+\alpha)} \right\} \\ \times \left\{ \frac{\theta\left(q^\alpha \frac{(-qz)}{1-q}\right)}{\theta\left(\frac{-qz}{1-q}\right)} (1-q)^{-\alpha} \right\} {}_2\varphi_1\left(q^\alpha, 0; q^{\alpha+1-\beta}; q, \frac{q^{1-\alpha-\beta}(1-q)}{z}\right), \end{aligned}$$

we obtain the conclusion. □

Combining lemma 5 and lemma 6, we obtain the theorem 4.

Acknowledgements

The author would like to give heartful thanks to Professor Yousuke Ohyama who provided carefully considered feedback and many valuable comments.

References

[1] G. D. Birkhoff, The generalized Riemann problem for linear differential equations and the allied problems for linear difference and q-difference equations, Proc. Am. Acad. Arts and Sciences, **49** (1914), 521 – 568.

[2] G. Gasper and M. Rahman, Basic Hypergeometric Series, 2nd ed, Cambridge, 2004.

[3] W. Hahn, Beiträge zur Theorie der Heineschen Reihen. Die 24 Integrale der Hypergeometrischen q-Differenzengleichung. Das q-Analogon der Laplace-Transformation. Math. Nachr. **2**, (1949). 340–379.

[4] T. Morita, A connection formula of the Hahn-Exton q-Bessel Function, SIGMA,**7** (2011), 115, 11pp.

[5] Y. Ohyama, A unified approach to q-special functions of the Laplace type, `arXiv:1103.5232`.

[6] G. N. Watson, The continuation of functions defined by generalized hypergeometric series, *Trans. Camb. Phil. Soc.* **21** (1910), 281–299.

[7] C. Zhang, Remarks on some basic hypergeometric series, in "Theory and Applications of Special Functions", Springer (2005), 479–491.

[8] C. Zhang, Sur les fonctions q-Bessel de Jackson, J. Approx. Theory, **122** (2003), 208–223.

[9] C. Zhang, Une sommation discrè pour des équations aux q-différences linéaires et à coefficients, analytiques: théorie générale et exemples, in "Differential Equations and Stokes Phenomenon", World Scientific (2002), 309–329.